



MacConkey Agar with CV, NaCl and 0.15% Bile Salt. 100 g / 500 g

Used for the detection of coliforms and other enteric pathogens.

Product Presentation:

Cat No.	Product description	Pack Size
11130010100	MacConkey Agar with CV, NaCl and 0.15% Bile Salt.	100 Gram
11130010500	MacConkey Agar with CV, NaCl and 0.15% Bile Salt.	500 Gram

Principle

MacConkey agars are slightly selective and differential plating media mainly used for the detection and isolation of Gram-negative organisms from clinical, dairy, food, water, pharmaceutical and industrial sources. It is also recommended for the selection and recovery of the *Enterobacteriaceae* and related enteric Gram-negative bacilli. USP recommends this medium for use in the performance of Microbial Limit Tests. The original medium contains protein, bile salts, sodium chloride and two dyes. The selective action of this medium is attributed to crystal violet and bile salts, which are inhibitory to most species of Gram-positive bacteria. Gram-negative bacteria usually grow well on the medium and are differentiated by their ability to ferment lactose. Lactose-fermenting strains grow as red or pink colonies and may be surrounded by a zone of acid precipitated bile.

The red colour is due to production of acid from lactose, absorption of neutral red and a subsequent colour change of the dye when the pH of medium falls below 6.8. Lactose non-fermenting strains, such as *Shigella* and *Salmonella* are colourless, transparent and typically do not alter appearance of the medium. Peptone, Proteose Peptone are sources of nitrogen, carbon, long chain amino acids and other nutrients. Lactose is a fermentable carbohydrate. Sodium chloride maintains the osmotic equilibrium. Bile salts and crystal violet are selective agents that inhibit growth of Gram-positive organisms. Neutral red is the pH indicator dye.

Composition

Ingredients

Grams / Litre

Peptone	17.00
Proteose Peptone	03.00
Lactose	10.00
Bile Salts	1.50
Sodium Chloride	05.00
Neutral Red	0.030
Crystal Violet	0.001
Agar	13.50

Final pH (at 25°C) 7.1±0.2

*Formula adjusted, standardized to suit performance parameters

FACTORY & OFFICE

Plot No. D 76 , Five Star MIDC Area, Kagal.

Dist. Kolhapur -416216 (M.S.)India.

Email : oxalispvtltd@outlook.com

Telefax : 0231-2305072

Phone : 0231-2305062

Mobile : +91 8805867810



Type of specimen

Clinical samples - Faeces, Urine and Other Pathological Material
Food and Dairy samples, Water samples, Pharmaceutical samples.

Specimen Collection and Handling

Ensure that all samples are properly labeled. Follow appropriate techniques for handling samples as per established guidelines. Some samples may require special handling, such as immediate refrigeration or protection from light, follow the standard procedure. The samples must be stored and tested within the permissible time duration. After use, contaminated materials must be sterilized by autoclaving before discarding.

Directions

- ✓ Suspend 50.00 g of powder in 1000 mL distilled water.
- ✓ Mix thoroughly.
- ✓ Boil to dissolve the medium completely. Avoid Overheating.
- ✓ Sterilize by autoclaving 121°C for 15 minutes or as per validated cycle.

Storage and Stability

- ✓ Store Dehydrated culture media in cool, dry place at 10°C-30°C away from direct light.
- ✓ Store prepared medium at 2°C-8°C. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Quality Control

Dehydrated Appearance: Light pink coloured homogeneous, free flowing powder

Prepared Appearance: Orange Red coloured, slightly opalescent gel forms in petridishes.

Growth Promotion Test: Growth promotion is carried out in accordance with the harmonized method of USP/EP/JP/IP and growth is observed after an incubation at 30°C-35°C for 18 to 72 hours.

Cultural Response :

Organism	Type Culture	Growth	Colour of Colony	Incubation Temperature	Incubation Period
<i>Escherichia coli</i>	ATCC 25922	Good	Pink with bile precipitate	30°C -35°C	18 Hours
<i>Salmonella enterica</i> <i>subsp. enterica serovar</i> <i>Typhimurium</i>	ATCC 14028	Good	Colourless	30°C -35°C	18 Hours

Inhibitory :

Organism	Type Culture	Growth	Incubation Temperature	Incubation Period
<i>Staphylococcus aureus</i>	ATCC 25923	Inhibited	30°C -35°C	48 Hours

FACTORY & OFFICE

Plot No. D 76 , Five Star MIDC Area, Kagal.
Dist. Kolhapur -416216 (M.S.)India.
Email : oxalispvtltd@outlook.com

Telefax : 0231-2305072
Phone : 0231-2305062
Mobile : +91 8805867810



Interpretation of Results

- ✓ Examination of plates for growth after completion of incubation period.

Warranty

- ✓ This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

Disposal

Disposal of infectious material and material that comes in to contact with clinical sample must be decontaminated and dispose of by autoclaving or incineration or established laboratory procedures. User must ensure safe disposal of used or unusable preparation of the products.

Reference

1. Murray P. R, Baron E, J., Jorgensen J. H., Pfaller M. A., Tenover F. C., Tenover F. C., (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.
2. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C

FACTORY & OFFICE

Plot No. D 76 , Five Star MIDC Area, Kagal.
Dist. Kolhapur -416216 (M.S.)India.
Email : oxalispvtltd@outlook.com

Telefax : 0231-2305072
Phone : 0231-2305062
Mobile : +91 8805867810